

The Role of Androgen Receptor Variant AR-V9 in Prostate Cancer

Yeung Ho¹, Jamie L Van Etten¹, Christine Henzler², Rendong Yang², Yingming Li¹, Elizabeth Tseng³, Ting Hon³, Tyson Clark³, and Scott M Dehm^{1,4}

UNIVERSITY OF MINNESOTA
Driven to DiscoverSM

¹Masonic Cancer Center, The University of Minnesota, Minneapolis, MN
²Minnesota Supercomputing Institute, The University of Minnesota, Minneapolis, MN
³Pacific Biosciences, Menlo Park, CA

⁴Department of Laboratory Medicine and Pathology, The University of Minnesota, Minneapolis, MN



ABSTRACT

The expression of androgen receptor (AR) variants is a frequent, yet poorly-understood mechanism of clinical resistance to AR-targeted therapy for castration-resistant prostate cancer (CRPC). Among the multiple AR variants expressed in CRPC, AR-V7 is considered the most clinically-relevant AR variant due to broad expression in CRPC, correlations of AR-V7 expression with clinical resistance, and growth inhibition when AR-V7 is knocked down in CRPC models. Therefore, efforts are under way to develop strategies for monitoring and inhibiting AR-V7 in castration-resistant prostate cancer (CRPC).

The aim of this study was to understand whether other AR variants are co-expressed with AR-V7 and promote resistance to AR-targeted therapies. To test this, we utilized RNA-seq to characterize AR expression in CRPC models. RNA-seq revealed the frequent coexpression of AR-V9 and AR-V7 in multiple CRPC models and metastases. Furthermore, long-read single-molecule real-time (SMRT) sequencing of AR isoforms revealed that AR-V7 and AR-V9 shared a common 3' terminal cryptic exon. To test this, we knocked down AR-V7 in prostate cancer cell lines and confirmed that AR-V9 mRNA and protein expression were also impacted. In reporter assays with AR-responsive promoters, AR-V9 functioned as a constitutive activator of androgen/AR signaling. Similarly, infection of AR-V9 lentiviral construct in LNCaP cells induced androgen-independent cell proliferation. In conclusion, these data implicate co-expression of AR-V9 with AR-V7 as an important component of constitutive AR signaling and therapeutic resistance in CRPC.

BACKGROUND

- Castration resistant prostate cancer (CRPC) develops when patients acquire resistance to androgen deprivation therapies (ADT).
- Expression of AR variants represents an important mechanism of the resistance to ADT.
- AR-V7 is considered the most clinically-relevant AR-V as the expression of AR-V7 in circulating tumor cells is associated with resistance to ADT in CRPC patients.
- It is not known whether other AR-Vs expressed in prostate cancer contribute to ADT resistance.

FIG 1. Frequent coexpression of AR-V9 and AR-V7 in CRPC models




Fig 1. RNA-seq reveals frequent coexpression of AR-V9 and AR-V7 in prostate cancer. A and C. Illumina RNA-seq read coverage along discrete regions of the AR gene in 22Rv1 cell line (A) and CWR-R1 cell line (C). B. Schematic depicting the relationship between AR and AR-V mRNA species and modular domains in AR and AR-V proteins. D. RNA-seq read coverage within intron 3 of the AR gene in the LuCaP 147 PDX and a liver metastasis biopsy.

FIG 2. AR-V7 and AR-V9 share a common 3' terminal cryptic exon




Fig 2. SMRT sequencing of AR isoforms in CRPC. A. Schematic of AR gene structure in 22Rv1 cells, which harbor a 35-kb AR intragenic tandem duplication. B. Exon composition and quantification of the 20 most abundant AR transcripts isolated from 22Rv1 cells by 3' rapid amplification of cDNA ends (RACE) using a forward primer anchored in AR exon 1. C. Exon composition and quantification of the 20 most abundant AR transcripts isolated from 22Rv1 cells by 3' RACE using a forward primer anchored in AR exon 3.

FIG 3. AR-V9 protein expression is inhibited by RNAi targeted to exon CE3




Fig 3. AR-V7 and AR-V9 contain overlap in 3'untranslated regions. A. Model for splicing of AR-V9 and AR-V7 derived from analysis of long-read sequence data. The model proposes that AR-V9 and AR-V7 originate from splicing of a common 3' terminal cryptic exon harboring two splice acceptor sites. B. siRNA targeting AR exon CE3 reduced mRNA expressions of both AR-V7 and AR-V9 in prostate cancer cell lines. In contrast, siRNA targeting AR exon CE5 reduced mRNA expression of AR-V9 but not AR-V7. C. Schematic of AR-V9 and AR-V7 proteins. D. siRNA targeting AR exon CE3 or CE5 reduced the protein expression level of AR-V9. In contrast, protein levels of AR-V7 was only reduced by siRNA targeting CE3.

FIG 4. AR-V9 is constitutively active and promotes androgen-independent growth




Fig 4. AR-V9 is constitutively active and promotes androgen-independent growth. A and B. AR-positive LNCaP cells were transfected with a PSA-driven luciferase reporter and luciferase assay was performed. C and D. AR-negative DU145 cells were transfected with an ARE-driven luciferase reporter. E and F. LNCaP cells were infected with GFP or AR-V9 lentivirus and assayed for proliferation by BrdU incorporation assay.

CONCLUSIONS

- AR-V9 was frequently coexpressed with AR-V7 in CRPC models.
- Both AR variant species were found to share a common 3' terminal cryptic exon.
- siRNA targeting AR exon CE3 reduced expression of both AR-V7 and AR-V9 mRNA and protein expression while siRNA targeting AR exon CE5 reduced expression of AR-V9 but not AR-V7.
- Similarly, siRNA targeting AR exon CE3 or CE5 reduced the protein expression level of AR-V9. In contrast, protein levels of AR-V7 were only reduced by siRNA targeting CE3.
- AR-V9 is constitutively active and promotes androgen-independent growth of CRPC cells.
- Co-expression of AR-V9 with AR-V7 may be an important component of constitutive AR signaling and therapeutic resistance in CRPC.

ACKNOWLEDGEMENTS

- ❖ This work was supported by a grant from the National Institutes of Health (R01 CA174777 to S.M.D.).
- ❖ We would like to thank all the people from Dehm lab for the help and support during the work.

Masonic Cancer Center
UNIVERSITY OF MINNESOTA
Comprehensive Cancer Center designated by the National Cancer Institute